

## ENTYVIO MEDICATION ORDER

Patient's Name (Last, First, Middle) \_\_\_\_\_ DOB: \_\_\_\_\_

■ **Diagnosis** Please specify the ICD10 code if using a diagnosis with a \_\_\_ noted.

K50.90 Ulcerative colitis

K50.9 \_\_\_\_\_ Crohn's disease (specific ICD10)

### ■ Details Needed for Authorization

- Proof of patient's negative latent TB test. If test is positive, proof that patient has begun therapy for latent TB.
- Is patient concurrently being treated with any other biologic? \_\_\_\_\_
- Does the patient have an intolerance, contraindication or hypersensitivity to any of the following agents, or has tried and failed on at least one with at least 3 months of therapy? If yes, circle all that apply. They are: 6-mercaptopurine, aminosaliclates, azathioprine, corticosteroids, mesalamine, methotrexate, sulfasalazine, hydroxychloroquine, Otezla, NSAIDs and leflunomide.
- Please provide documented failure, contraindication, or ineffective response at maximum tolerated doses to a minimum (3) month trial on previous therapy with a TNF modifier such as Humira, Simponi, or infliximab (Avsola, Inflectra, Remicade or Renflexis).

### ■ Entyvio (vedolizumab) Medication Order

Patient's height in ft/in: \_\_\_\_\_ Patient's weight in lbs: \_\_\_\_\_

Select all doses required.

Starting dose: 300mg in 250ml normal saline over about 30 minutes at weeks 0, 2 and 6.

Maintenance dose: 300mg in 250ml normal saline over about 30 minutes every 8 weeks for \_\_\_\_\_ months.

Post infusion flush with normal saline. Check vitals and monitor for signs and symptoms at start, throughout infusion, and after completion.

### ■ Rescue Management in case of Infusion Therapy Reaction

These include fever, chills, rigors, headache, rash, itching, swelling, edema, nausea, vomiting, abdominal pain, hypotension, and respiratory distress.

- Stop medication infusion and start normal saline infusion at 50 ml/hr. Call ordering provider to report reaction.
- Follow standing reaction orders, including diphenhydramine, methylprednisolone, albuterol and oxygen as needed.
- For severe reactions, administer Epi-pen or equivalent and call 911. Repeat if severe symptoms persist.

### ■ Ordering Provider Authorization

Provider's Signature: \_\_\_\_\_ Name: \_\_\_\_\_ Date: \_\_\_\_\_

Address: \_\_\_\_\_

Phone: \_\_\_\_\_ Fax: \_\_\_\_\_ NPI #: \_\_\_\_\_ License: \_\_\_\_\_

### STANDARD DOCUMENTATION TO INCLUDE:

- Patient demographics and insurance, including card scans (both medical and pharmacy benefit cards, both sides).
- Most recent chart notes and, if available, last history and physical. All relevant scans, tests and laboratory results.
- If new medication for patient, chart notes which include decision to begin treatment. If not, provide last treatment date.

**Fax this order and supporting documentation to (732) 329-2322.**